

A COUNTEREXAMPLE TO THE LOCAL-GLOBAL PRINCIPLE OF LINEAR DEPENDENCE FOR ABELIAN VARIETIES

PETER JOSSEN AND ANTONELLA PERUCCA

ABSTRACT. Let A be an abelian variety defined over a number field k . Let P be a point in $A(k)$ and let X be a subgroup of $A(k)$. Gajda in 2002 asked whether it is true that the point P belongs to X if and only if the point $(P \bmod \mathfrak{p})$ belongs to $(X \bmod \mathfrak{p})$ for all but finitely many primes $\mathfrak{p}$ of k . We answer negatively to Gajda's question.

Let A be an abelian variety defined over a number field k . Let P be a point in $A(k)$ and let X be a subgroup of $A(k)$. Suppose that for all but finitely many primes $\mathfrak{p}$ of k the point $(P \bmod \mathfrak{p})$ belongs to $(X \bmod \mathfrak{p})$. Is it true that P belongs to X ? This question, which was formulated by Gajda in 2002, was named the problem of detecting linear dependence. The problem was addressed in several papers (see the bibliography) but the question was still open. In this note we show that the answer to Gajda's question is negative by providing a counterexample.

Let k be a number field and let E be an elliptic curve without complex multiplication over k such that there are points P_1, P_2, P_3 in $E(k)$ which are $\mathbb{Z}$ -linearly independent. Let $P \in E^3(k)$ and $X \subseteq E^3(k)$ be the following:

$$P := \begin{pmatrix} P_1 \\ P_2 \\ P_3 \end{pmatrix} \quad X := \left\{ MP \in E^3(k) \mid M \in \text{Mat}(3, \mathbb{Z}), \text{tr } M = 0 \right\}$$

So the group X consists of the images of the point P via the endomorphisms of E^3 with trace zero. Since the points P_i are $\mathbb{Z}$ -independent and since E has no complex multiplication, the point P does not belong to X . Notice that no non-zero multiple of P belongs to X .

Claim. *Let $\mathfrak{p}$ be a prime of k where E has good reduction. The image of P under the reduction map modulo $\mathfrak{p}$ belongs to the image of X .*

For the rest of this note, we fix a prime $\mathfrak{p}$ of good reduction for E . We write κ for the residue field of k at $\mathfrak{p}$. To ease notation, we now let E denote the reduction of the given elliptic curve modulo $\mathfrak{p}$ and write P_1, P_2, P_3, P and X for the image of the given points and the given subgroup under the reduction map modulo $\mathfrak{p}$.

Our aim is to find an integer matrix M of trace zero such that $P = MP$ in $E^3(\kappa)$.

Let α_1 be the smallest positive integer such that $\alpha_1 P_1$ is a linear combination of P_2 and P_3 . Similarly define α_2 and α_3 for P_2 and P_3 respectively. There are integers m_{ij} such that

$$\begin{aligned}\alpha_1 P_1 + m_{12} P_2 + m_{13} P_3 &= O \\ m_{21} P_1 + \alpha_2 P_2 + m_{23} P_3 &= O \\ m_{31} P_1 + m_{32} P_2 + \alpha_3 P_3 &= O\end{aligned}$$

Assume that the greatest common divisor of α_1 , α_2 and α_3 is 1 (we prove this assumption later). We can thus find integers a_1, a_2, a_3 such that

$$3 = \alpha_1 a_1 + \alpha_2 a_2 + \alpha_3 a_3$$

Write $m_{ii} := 1 - \alpha_i a_i$, so that in particular $m_{11} + m_{22} + m_{33} = 0$. Then we have

$$\begin{pmatrix} m_{11} & -a_1 m_{12} & -a_1 m_{13} \\ -a_2 m_{21} & m_{22} & -a_2 m_{23} \\ -a_3 m_{31} & -a_3 m_{32} & m_{33} \end{pmatrix} \begin{pmatrix} P_1 \\ P_2 \\ P_3 \end{pmatrix} = \begin{pmatrix} P_1 \\ P_2 \\ P_3 \end{pmatrix}$$

Notice that the above matrix has integer entries and trace zero. Hence we are left to prove that the greatest common divisor of α_1 , α_2 and α_3 is indeed 1.

Fix a prime number ℓ and let us show that ℓ does not divide $\gcd(\alpha_1, \alpha_2, \alpha_3)$. Suppose on the contrary that ℓ divides $\gcd(\alpha_1, \alpha_2, \alpha_3)$. This is equivalent to saying that ℓ divides all the coefficients appearing in any linear relation between P_1 , P_2 and P_3 . In particular, this implies that ℓ divides the order of P_1 , P_2 and P_3 in $E(\kappa)$.

It is well-known that the group $E(\kappa)[\ell]$ is either trivial, isomorphic to $\mathbb{Z}/\ell\mathbb{Z}$ or isomorphic to $(\mathbb{Z}/\ell\mathbb{Z})^2$. In any case, the intersection $X \cap E(\kappa)[\ell]$ is generated by two elements or less. Without loss of generality, let us suppose that the subgroup of X generated by P_2 and P_3 contains $X \cap E(\kappa)[\ell]$.

We are supposing that ℓ divides all the coefficients appearing in any linear relation of the points. Let $\alpha_1 = x_1 \ell$ and write $x_1 \ell P_1 + x_2 \ell P_2 + x_3 \ell P_3 = O$ for some integers x_2 and x_3 . It follows that

$$x_1 P_1 + x_2 P_2 + x_3 P_3 = T$$

for some point T in $X \cap E(\kappa)[\ell]$. The point T is a linear combination of P_2 and P_3 . This contradicts the minimality of α_1 .

REFERENCES

1. G. Banaszak, *On a Hasse principle for Mordell-Weil groups*, C. R. Math. Acad. Sci. Paris. (to appear) (2008), 1–8.
2. G. Banaszak, W. Gajda, and P. Krasoń, *Detecting linear dependence by reduction maps*, J. Number Theory **115** (2005), no. 2, 322–342.
3. G. Banaszak and P. Krasoń, *On arithmetic in Mordell-Weil groups*, preprint, arXiv:math/0904.2848 (2009), 1–17.
4. S. Barańczuk, *On generalization of the support problem of Erdős and its analogues for abelian varieties and K-theory*, preprint, arXiv:math/0809.1991 (2008), 1–6.
5. W. Gajda and K. Górniewicz, *Linear dependence in Mordell-Weil groups*, J. Reine Angew. Math. (2009), 1–15.
6. P. Jossen, *Detecting linear dependence on a semiabelian variety*, preprint, arXiv:math/0903.5271 (2009), 1–16.

7. C. Khare, *Compatible systems of mod p Galois representations and Hecke characters*, Math. Res. Lett. **10** (2003), 71–83.
8. E. Kowalski, *Some local–global applications of Kummer theory*, Manuscripta Math. **111** (2003), no. 1, 105–139.
9. A. Perucca, *On the problem of detecting linear dependence for products of abelian varieties and tori*, preprint, arXiv:math/0811.1495 (2008), 1–9.
10. A. Schinzel, *On power residues and exponential congruences*, Acta Arith. **27** (1975), 397–420.
11. T. Weston, *Kummer theory of abelian varieties and reduction of Mordell–Weil groups*, Acta Arith. **110** (2003), 77–88.

Peter Jossen

Department of Mathematics, Central European University

Nádor u. 9, 1051 Budapest, Hungary

`peter.jossen@gmail.com`

Antonella Perucca

Section des Mathématiques, École Polytechnique Fédérale de Lausanne

EPFL, MA C3 604, Station 8, 1015 Lausanne, Switzerland

`antonella.perucca@epfl.ch`